

A highly stereoselective route to medium-ring-sized trans-alkenolides via oxidative fragmentation of bicyclic oxycyclopropane precursors : application to the synthesis of (+)-recifeiolide
Zubrytski, Dzmitry; Kananovich, Dzmitry; Kulinkovich, Oleg Tetrahedron 2014 / p. 2944-2950 : ill
<https://www.sciencedirect.com/science/article/abs/pii/S0040402014003524>

Horner-Wadsworth-Emmons kondensatsiooni intermediaatide kvantkeemiline uurimine
Martin, Ivar; Vares, L.; Rein, T. XXIII Eesti keemiapäevad : teaduskonverentsi ettekannete referaadid 1997 / lk. 80

Organocatalytic stereoselective [8+2] and [6+4] cycloadditions

Mose, Rasmus; **Preegel, Gert**; Larsen, Jesper; Jakobsen, Sofie; Iversen, Eva Hogh; Jorgensen, Karl Anker Nature chemistry 2017 / p. 487-492 : ill <http://dx.doi.org/10.1038/nchem.2682>

Quantum chemical exploration of intermediates in the Horner-Wadsworth-Emmons condensation

Martin, Ivar; Vares, L.; Rein, T. 23rd Estonian Chemistry Days : abstracts of scientific conference 1997 / p. 89

Regio- and stereoselective synthesis of chiral dihydroxyketones

Nidu, Allan; Paju, Anne; Eek, Margus; Kanger, Tõnis; Pehk, Tõnis; Lopp, Margus Komppa Centenary Symposium : July 6-9, 2003, Espoo, Finland : abstracts 2003 / p. A70

Titanacyclopropane reagents for stereoselective organic synthesis

Kulinkovich, Oleg BOS 2012 : International Conference on Organic Synthesis : July 1-4, 2012, Tallinn, Estonia : program and abstracts 2012 / p. 22