

Activation of 11R-lipoxygenase is fully Ca²⁺-dependent and controlled by the phospholipid composition of the target membrane

Järving, Reet; Lõokene, Aivar; Kurg, Reet; **Siimon, Liina; Järving, Ivar; Samel, Nigulas** Biochemistry 2012 / p. 3310-3320 : ill
<https://pubs.acs.org/doi/10.1021/bi201690z>

Apolipoprotein A-V interaction with members of the low density lipoprotein receptor gene family

Nilsson, Stefan; **Lõokene, Aivar;** Beckstead, Jennifer; Gliemann, Jorgen; Ryan, Robert; Olivecrona, Gunilla Biochemistry 2007 / 12, p. 3896-3904 : ill
<https://pubs.acs.org/doi/10.1021/bi7000533#:~:text=Experiments%20using%20surface%20plasmon%20resonance,low%20density%20lipoprotein%20receptor%20family.>

Brain-specific metallothionein-3 has higher metal-binding capacity than ubiquitous metallothioneins and binds metals noncooperatively

Palumaa, Peep; Eriste, Elo; **Njunkova, Olga;** Pokras, Lesja; Jorvall, H.; Sillard, Rannar Biochemistry 2002 / 19, p.6158-6163
<https://pubmed.ncbi.nlm.nih.gov/11994011/>

Characterization of protein kinase ULK3 regulation by phosphorylation and inhibition by small molecule SU6668

Kasak, Lagle; Näks, Mihkel; Eek, Priit; Piirsoo, Alla; **Bhadoria, Rohit; Starkov, Pavel;** Saarma, Merilin; Kasvandik, Sergo; **Piirsoo, Marko** Biochemistry 2018 / p. 5456–5465 <https://doi.org/10.1021/acs.biochem.8b00356> [Journal metrics at Scopus](#) [Article at Scopus](#) [Journal metrics at WOS](#) [Article at WOS](#)

Redox and metal ion binding properties of human insulin-like growth factor 1 determined by electrospray ionization mass spectrometry

Smirnova, Julia; Muhhina, Jekaterina; Tõugu, Vello; Palumaa, Peep Biochemistry 2012 / p. 5851-5859 : ill
<https://pubs.acs.org/doi/10.1021/bi300494s>