

Accessibility of published QSAR models : ecotoxic endpoints

Kahn, Iiris; Ahte, Priit; Sild, Sulev; Maran, Uko ALTEX proceedings. Vol. 4, No. 2 2015 / p. 119

Accessibility of published QSAR models : environmental fate endpoints

Kahn, Iiris; Ahte, Priit; Piir, Geven; Sild, Sulev; Maran, Uko Conferentia Chemometrica 2015 : Budapest, MTA TTK, September 13-16, 2015 2015 / p. P33

Best practices for QSAR model reporting: physical and chemical properties, ecotoxicity, environmental fate, human health, and toxicokinetics endpoints

Piir, Geven; Kahn, Iiris; García-Sosa, Alfonso T.; Sild, Sulev; Ahte, Priit; Maran, Uko Environmental Health Perspectives 2018 / art. 126001 ; 20 p.: ill <https://doi.org/10.1289/EHP3264> [Journal metrics at Scopus](#) [Article at Scopus](#) [Journal metrics at WOS](#) [Article at WOS](#)

Comparative quantitative structure-activity-activity relationships for toxicity to Tetrahymena pyriformis and Pimephales promelas

Kahn, Iiris; Maran, Uko; Benfenati, Emilio; Netzeva, Tatiana; Schults, T.Wayne; Cronin, Mark ATLA = Alternatives to laboratory animals 2007 / 1, p. 15-24 https://www.researchgate.net/publication/6411335_Comparative_Quantitative_Structure-Activity-Activity_Relationships_for_Toxicity_to_Tetrahymena_pyriformis_and_Pimephales_promelas

Improved efficiency of focal point conformational analysis with truncated correlation consistent basis sets

Kahn, Kalju; Kahn, Iiris Journal of computational chemistry 2008 / 6, p. 900-911 <https://onlinelibrary.wiley.com/doi/abs/10.1002/jcc.20848>

In silico approach to finding new scaffolds for LRRK2 inhibition

Kahn, Iiris; Lomaka, Andre; Karelson, Mati European Pharma Summit : Berlin, Germany, May 5-8, 2015 : abstracts 2015 / [1] p

Modeling the toxicity of chemicals to Tetrahymena pyriformis using heuristic multilinear regression and heuristic back-propagation neural networks

Kahn, Iiris; Sild, Sulev; Maran, Uko Journal of chemical information and modeling 2007 / 6, p. 2271-2279 <https://pubs.acs.org/doi/10.1021/ci700231c>

QSPR treatment of the soil sorption coefficients of organic pollutants

Kahn, Iiris; Fara, Dan; Karelson, Mati; Maran, Uko; Andersson, Patrik L. Journal of chemical information and modeling 2005 / 1, p. 94-105 <https://pubs.acs.org/doi/10.1021/ci0498766>

Quantitative structure-activity relationship (QSAR) modeling of EC50 of aquatic toxicities for Daphnia magna

Katritzky, Alan R.; Slavov, Svetoslav H.; Stoyanova-Slavova, Iva B.; Kahn, Iiris; Karelson, Mati Journal of toxicology and environmental health. Part A, Current issues 2009 / 19, p. 1181-1190 : ill <https://pubmed.ncbi.nlm.nih.gov/20077186/>

Quantitative structure-activity relationships of environmentally relevant properties

Kahn, Iiris 2007

Reproducibility of QSAR/QSPR models in the scientific literature - perspective for regulatory use

Kahn, Iiris; Piir, Geven; Oja, Mare Scientific programme & book of abstracts : QSAR 2014 2014 / p. 109

Topological fingerprints as an aid in finding structural patterns for LRRK2 inhibition

Kahn, Iiris; Lomaka, Andre; Karelson, Mati Molecular informatics 2014 / p. 269-275 : ill <https://doi.org/10.1002/minf.201300057> [Journal metrics at Scopus](#) [Article at Scopus](#) [Journal metrics at WOS](#) [Article at WOS](#)

Unsubstituted oxalalix[n]arenes (n=4 and 8): a conformational study in solution and solid state and interaction studies with aromatic guests

Peterson, Anna; Kaabel, Sandra; Kahn, Iiris; Pehk, Tõnis; Aav, Riina; Adamson, Jasper ChemistrySelect 2018 / p. 9091 - 9095 <https://doi.org/10.1002/slct.201801590> [Journal metrics at Scopus](#) [Article at Scopus](#) [Journal metrics at WOS](#) [Article at WOS](#)