

Abiotic factors controlling the cyanobacterial bloom occurrence in the Gulf of Finland

Lips, Inga 2005 <http://dspace.ut.ee/handle/10062/940>

A comparative SEM study of ossicles in the Pleuronectiformes (Teleostei) of the Baltic Sea

Märss, Tiiu; Wilson, Mark; Lees, Janek; Saat, Toomas; Špilev, Heli Proceedings of the Estonian Academy of Sciences 2015 / p. 495-517 : ill <https://doi.org/10.3176/proc.2015.4.05> Journal metrics at Scopus Article at Scopus Journal metrics at WOS Article at WOS

Life history of anadromous burbot (*Lota lota*, Linneaus) in the brackish Baltic Sea inferred from otolith microchemistry

Rohtla, Mehis; Vetemaa, Markus; Taal, Imre; Svärdsden, Roland; Urtson, Kristjan; Saks, Lauri; Verliin, Aare; Kesler, Martin; Saat, Toomas Ecology of freshwater fish 2014 / p. 141-148 : ill <https://doi.org/10.1111/eff.12057> Journal metrics at Scopus Article at Scopus Journal metrics at WOS Article at WOS

The morphology and sculpture of ossicles in the Cottidae (Teleostei) of the Baltic Sea

Märss, Tiiu; Lees, Janek; Wilson, Mark V.H.; Saat, Toomas; Špilev, Heli Estonian journal of earth sciences 2010 / 3, p. 216-237 : ill https://artiklid.elnet.ee/record=b2163755*est

The morphology and sculpture of ossicles in the Cyclopteridae and Liparidae (Teleostei) of the Baltic Sea

Märss, Tiiu; Lees, Janek; Wilson, Mark; Saat, Toomas; Špilev, Heli Estonian journal of earth sciences 2010 / p. 263-276 : ill https://artiklid.elnet.ee/record=b2183599*est