

Functional diversity of human basic helix-loop-helix transcription factor TCF4 isoforms generated by alternative 5' exon usage and splicing

Sepp, Mari; Kannike, Kaja; Eesmaa, Ave; Urb, Mari; Timmus, Tõnis PLoS ONE 2011 / p. e22138 [14 p.] : ill

<https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0022138>

Internalisation of cell-penetrating peptides into tobacco protoplasts

Mäe, Maarja; Myrberg, Helena; Jiang, Yang; Paves, Heiti; Valkna, Andres; Langel, Ülo Biochimica et biophysica acta : biomembrans 2005 / 2, p. 101-107 : ill <https://pubmed.ncbi.nlm.nih.gov/15893512/>

Refinement of a quantitative structure–activity relationship model for prediction of cell-penetrating peptide based transfection systems

Dowaidar, Moataz; Regberg, Jakob; **Dobchev, Dimitar Atanasov**; Lehto, Tõnis; Hällbrink, Mattias; **Karelson, Mati**; Langel, Ülo International journal of peptide research and therapeutics 2017 / p. 91-100 : ill <https://doi.org/10.1007/s10989-016-9542-8> [Journal metrics at Scopus](#) [Article at Scopus](#) [Journal metrics at WOS](#) [Article at WOS](#)