

Forkhead transcription factor FOXO3a levels are increased in Huntington disease because of overactivated positive autoregulatory loop

Kannike, Kaja; Sepp, Mari; Zuccato, Chiara; Cattaneo, Elena; Timmus, Tõnis Journal of biological chemistry 2014 / p. 32845-32857 : ill <https://doi.org/10.1074/jbc.M114.612424> [Journal metrics at Scopus](#) [Article at Scopus](#) [Journal metrics at WOS](#) [Article at WOS](#)

Functional diversity of human basic helix-loop-helix transcription factor TCF4 isoforms generated by alternative 5' exon usage and splicing

Sepp, Mari; Kannike, Kaja; Eesmaa, Ave; Urb, Mari; Timmus, Tõnis PLoS ONE 2011 / p. e22138 [14 p.] : ill <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0022138>

Partial deletion of TCF4 in three generation family with nonsyndromic intellectual disability, without features of Pitt-Hopkins syndrome

Kharbanda, Mira; Kannike, Kaja; Lampe, Anne; Berg, Jonathan; Timmus, Tõnis; Sepp, Mari European journal of medical genetics 2016 / p. 310-314 : ill <https://doi.org/10.1016/j.ejmg.2016.04.003> [Journal metrics at Scopus](#) [Article at Scopus](#) [Journal metrics at WOS](#) [Article at WOS](#)