

Cloning and expression of ATP N-glycosidase from the freshwater sponge *Ephydatia muelleri*

Reintamm, Tõnu; Vallmann, Kerli; Kolk, Kaidi; Päre, Mailis; Lopp, Annika; Aas-Valleriani, Nele; Kelve, Merike Biochimie 2019 / p. 126-129 <https://doi.org/10.1016/j.biochi.2018.12.018> [Journal metrics at Scopus](#) [Article at Scopus](#) [Journal metrics at WOS](#) [Article at WOS](#)

Enzymatically active 2',5'-oligoadenylate synthetases are widely distributed among Metazoa, including protostome lineage

Päre, Mailis; Kuusksalu, Anne; Lopp, Annika; Hansen Kjaer, Karina; Justesen, Just; Kelve, Merike Biochimie 2014 / p. 200-209 : ill <https://doi.org/10.1016/j.biochi.2013.10.015> [Journal metrics at Scopus](#) [Article at Scopus](#) [Journal metrics at WOS](#) [Article at WOS](#)

A novel endoribonuclease from the marine sponge *Tethya aurantium* specific to 2',5'-phosphodiester bonds

Lopp, Margus; Reintamm, Tõnu; Kuusksalu, Anne; Rosa, Salvatore, de; Kelve, Merike Biochimie 2012 / p. 1635-1646 : ill <https://doi.org/10.1016/j.biochi.2012.04.002> <https://www.sciencedirect.com/science/article/pii/S0300908412001368>