

Cell type-specific labelling of newly synthesized proteins by puromycin inactivation

Cabrera-Cabrera, Florencia; Tull, Helena; Capuana, Roberta; Kasvandik, Sergio; **Timmusk, Tõnis; Koppel, Indrek** Journal of biological chemistry 2023 / art. 105129 <https://doi.org/10.1016/j.jbc.2023.105129>

Human glial cell line-derived neurotrophic factor receptor alpha4 is the receptor for persephin, and is predominantly expressed in normal and malignant thyroid medullary cells

Lindahl, Maria; Poteryaev, Dimitry; Yu, Liying; Arumäe, Urmas; Timmusk, Tõnis; Bongarzone, Italia; Aiello, Antonella; Pierotti, Marco A.; Airaksinen, Matti S.; **Saarma, Mart** Journal of biological chemistry 2001 / 12, p. 9344-9351 : ill
<https://doi.org/10.1074/jbc.M008279200>

Other neurotrophic factors : Glial cell line-derived neurotrophic factor

Saarma, Mart; Sariola, H. Microscopy research and technique 1999 / p. 292-302

Promotion of seminomatous tumors by targeted overexpression of glial cell line-derived neurotrophic factor in mouse testis

Meng, X.; Rooij, D.G. de; Westerdahl, K.; **Saarma, Mart;** Sariola, H. Cancer research 2001 / p. 3267-3271
<https://pubmed.ncbi.nlm.nih.gov/11309277/>

Stroke induces widespread changes of gene expression for glial cell line-derived neurotrophic factor family receptors in the adult rat brain

Arvidsson, A.; Kokaia, Zaal; Airaksinen, Matti S.; **Saarma, Mart;** Lindvall, O. Neuroscience 2001 / p. 27-41
<https://pubmed.ncbi.nlm.nih.gov/11564414/>