Arabidopsis thaliana myosin XIK is involved in root hair as well as trichome morphogenesis on stems and leaves Ojangu, Eve-Ly; Järve, Kristel; Paves, Heiti; Truve, Erkki Protoplasma 2007 / 3/4, p. 193-202 https://link.springer.com/article/10.1007/s00709-006-0233-8

Comparative analysis of the composition of the essential oil from the shoots, leaves and stems the wild Ledum palustre L. from Estonia

Gretšušnikova, Tatjana; Järvan, Kerly; Orav, Anne; Koel, Mihkel Procedia chemistry 2010 / p. 168-173

Comparison of essential oils composition from shoots, leaves and stems of the wild Ledum palustre from Estonia Gretšušnikova, Tatjana; Orav, Anne; Koel, Mihkel; Järvan, Kerly NoSSS2009: 5th Conference on Separation and Related Techniques by Nordic Separation Science Society: 26-29 August, 2009, Tallinn University of Technology, Estonia: abstract book and program 2009 / p. 90

Identification of bioactive compounds in leaves and stems of Aegopodium podagraria by various analytical technics Orav, Anne; Viitak, Anu; Helmja, Kati; Vaher, Merike NoSSS2009: 5th Conference on Separation and Related Techniques by Nordic Separation Science Society: 26-29 August, 2009, Tallinn University of Technology, Estonia: abstract book and program 2009 / p. 144

Identification of bioactive compounds in the leaves and stems of Aegopodium podagraria by various analytical techniques

Orav, Anne; Viitak, Anu; Vaher, Merike Procedia chemistry 2010 / p. 152-160 : ill