Behaviour mechanisms and correlation between lead (Pb) and its isotope 210Pb in industrial residue as an indicator for waste characterization
Vaasma, Taavi; **Bitjukova, Liidia**; Kiisk, Madis; Özden, Banu; Tkaczyk, Alan Henry Environmental technology 2016 / p. 3208-3218 : ill http://dx.doi.org/10.1080/09593330.2016.1181673

Long-term modelling of fly ash and radionuclide emissions as well as deposition fluxes due to the operation of large oil shale-fired power plants
Vaasma, Taavi; Kaasik, Marko; **Loosaar, Jüri**; Kiisk, Madis; Tkaczyk, Alan Henry Journal of environmental radioactivity 2017 / p. 232-244 : ill https://doi.org/10.1016/j.jenvrad.2017.08.017

Pb-210 and Po-210 atmospheric releases via fly ash from oil shale-fired power plants
Vaasma, Taavi; **Loosaar, Jüri**; Gyakwaa, Francis; Kiisk, Madis; Özden, Banu; Tkaczyk, Alan Henry Environmental pollution 2017 / p. 210-218 : ill https://doi.org/10.1016/j.envpol.2016.12.054

Radionuclide concentration variations in residues from potential NORM industries in Estonia : the example of oil shale-fired power plants
Vaasma, Taavi; Kiisk, Madis; Realo, Enn; **Loosaar, Jüri**; Tkaczyk, Alan Henry V. Terrestrial Radioisotopes in Environment : International Conference on Environmental Protection 2016 / p. 30 http://dx.doi.org/10.18428/TREICEP-2016

Radionuclide concentration variations in the fuel and residues of oil shale-fired power plants : estimations of the radiological characteristics over a 2-year period
Vaasma, Taavi; **Loosaar, Jüri**; Kiisk, Madis; Tkaczyk, Alan Henry Journal of environmental radioactivity 2017 / p. 25-33 : ill https://doi.org/10.1016/j.jenvrad.2016.10.005