

Distribution of GABA receptor subunit transcripts in the rat brain

Wegelius, K.; Pasternack, M.; Hiltunen, J.O.; Rivera, Claudio; Kaila, Kai; **Saarma, Mart**; Reeben, M. European journal of neuroscience 1998 / p. 350-357

GDNF family receptors in the embryonic and postnatal rat heart and reduced cholinergic innervation in mice hearts lacking Ret or GFRa2

Hiltunen, J.O.; Laurikainen, A.; Airaksinen, Matti S.; **Saarma, Mart** Developmental dynamics 2000 / p. 28-39
<https://pubmed.ncbi.nlm.nih.gov/10974669/>

Glial cell line-derived neurotrophic factor is expressed in penis of adult rat and retrogradely transported in penile parasympathetic and sensory nerves

Laurikainen, A.; Hiltunen, J.O.; Vanhatalo, S.; Klinge, E.; **Saarma, Mart** Cell and tissue research 2000 / p. 321-329
<https://pubmed.ncbi.nlm.nih.gov/11151444/>

Nerve growth factor and brain-derived neurotrophic factor mRNAs are regulated in distinct cell populations of rat heart after ischaemia and reperfusion

Hiltunen, J.O.; Laurikainen, A.; Väkeva, A.; Meri, S.; **Saarma, Mart** Journal of pathology 2001 / p. 247-253
<https://pubmed.ncbi.nlm.nih.gov/11400155/>

Neurotrophin-3 is a target-derived neurotrophic factor for penile erection-inducing neurons

Hiltunen, J.O.; Laurikainen, A.; Klinge, E.; **Saarma, Mart** Neuroscience 2005 / 1, p. 51-58
<https://www.sciencedirect.com/science/article/pii/S0306452205000965>

Neurturin is a neurotrophic factor for penile parasympathetic neurons in adult rat

Laurikainen, A.; Hiltunen, J.O.; Thomas-Crusells, J.; Vanhatalo, S.; Arumäe, Urmas; Airaksinen, Matti S.; Klinge, E.; **Saarma, Mart** Journal of neurobiology 2000 / p. 198-205 <https://pubmed.ncbi.nlm.nih.gov/10770848/>

The first cysteine-rich domain of the receptor GFR[alpha]1 stabilizes the binding of GDNF

Virtanen, Heidi; Yang, Jianmin; Bessalov, Maxim M.; Hiltunen, J.O.; Leppänen, V.M.; Kalkkinen, Nisse; Goldman, A.; **Saarma, Mart**; Runeberg-Roos, Pia Biochemical journal 2005 / 3, p. 817-824 <https://pubmed.ncbi.nlm.nih.gov/15610063/>

The messenger RNAs for both glial cell line-derived neurotrophic factor receptors, C-ret and GDNFR, are induced in the rat brain in response to Kainate-induced excitation

Reeben, M.; Laurikainen, A.; Hiltunen, J.O.; Castren, Eero; **Saarma, Mart** Neuroscience 1998 / p. 151-159

Towards understanding the neuronal pathogenesis of aspartylglycosaminuria : expression of aspartylglycosaminidase in brain during development

Uusitalo, A.; Tenhunen, K.; Heinonen, O.; Hiltunen, J.O.; **Saarma, Mart**; Haltia, M.; Jalanko, A.; Palotie, L. Molecular genetics development 1999 / p. 294-307