

GDNF family ligands and receptors are differentially regulated after brain insults in the rat

Kokaia, Zaal; Airaksinen, Matti S.; Nanobashvili, A.; Larsson, E.; Kujamäki, E.; Lindvall, O.; **Saarma, Mart** European journal of neuroscience 1999 / p. 1202-1216

MANF is widely expressed in mammalian tissues and differently regulated after ischemic and epileptic insults in rodent brain

Lindholm, Päivi; Peränen, Johan; Andressoo, Jaan-Olle; Kalkkinen, Nisse; Kokaia, Zaal; Lindvall, Olle; **Timmusk, Tõnis; Saarma, Mart** Molecular and cellular neuroscience 2008 / p. 356-371 : ill

Seizures induce widespread upregulation of cystatin B, the gene mutated in progressive myoclonus epilepsy, in rat forebrain neurons

D'Amato, E.; Kokaia, Zaal; Nanobashvili, A.; Reeben, M.; Lehesjoki, A.-E.; **Saarma, Mart**; Lindvall, O. European journal of neuroscience 2000 / p. 1687-1695 <https://pubmed.ncbi.nlm.nih.gov/10792446/>

Stroke induces widespread changes of gene expression for glial cell line-derived neurotrophic factor family receptors in the adult rat brain

Arvidsson, A.; Kokaia, Zaal; Airaksinen, Matti S.; **Saarma, Mart**; Lindvall, O. Neuroscience 2001 / p. 27-41
<https://pubmed.ncbi.nlm.nih.gov/11564414/>

The neuronal ceroid lipofuscinosis Cln8 gene expression is developmentally regulated in mouse brain and up-regulated in the hippocampal kindling model of epilepsy

Lonka, Liina; Aalto, Antti; Kopra, Outi; Kuronen, Mervi; Kokaia, Zaal; **Saarma, Mart**; Lehesjoki, Anna-Elina BMC neuroscience 2005 / 1, p. 27 <https://doi.org/10.1186/1471-2202-6-27>