

Optical method for cardiovascular risk marker uric acid removal assessment during dialysis

Holmar, Jana; Fridolin, Ivo; Uhlin, Nils Fredrik Arne; Lauri, Kai; Luman, Merike The scientific world journal 2012 / p. 1-8 : ill

Optical method for uric acid removal assessment during dialysis = Optiline meetod kusihappe eemaldamise määramiseks dialüüsiravi käigus

Holmar, Jana 2013 https://www.esther.ee/record=b3004743*est

Paracetamol interferes uric acid levels in uraemic patients revealed by monitoring spent dialysate

Tanner, Risto; Arund, Jürgen; Fridolin, Ivo; Luman, Merike ISRN nephrology 2013 / p. 1-4 : ill

POS-648 medicines intake influences accuracy of the uremic retention molecules' optical monitoring in spent dialysate : the case of uremic toxin uric acid and paracetamol

Paats, Joosep; Adoberg, Annika; Arund, Jürgen; Fridolin, Ivo; Holmar, Jana; Leis, Liisi; Pilt, Kristjan; Tanner, Risto; Uhlin, Nils Fredrik Arne; Luman, Merike Kidney International Reports 2022 / p. S277-S278 : ill <https://doi.org/10.1016/j.kir.2022.01.681>

Treatment with paracetamol can interfere with the intradialytic optical estimation in spent dialysate of uric acid but not of indoxyl sulfate

Adoberg, Annika; Paats, Joosep; Arund, Jürgen; Dhondt, Annemieke; Fridolin, Ivo; Glorieux, Griet; Holmar, Jana; Lauri, Kai; Leis, Liisi; Luman, Merike; Pilt, Kristjan; Uhlin, Nils Fredrik Arne; Tanner, Risto Toxins 2022 / art. 610

<https://doi.org/10.3390/toxins14090610> Journal metrics at Scopus Article at Scopus Journal metrics at WOS Article at WOS