

Could urea in spent dialysate be a marker for monitoring removal of 4-pyridoxic acid?

Kalle, Sigrid; Tanner, Risto; Arund, Jürgen; Tomson, Ruth; Luman, Merike; Fridolin, Ivo XIII Baltic Nephrology Conference : October 13-15, 2016, Jūrmala, Latvia : final programme 2016 / p. 6 <http://nefrologs.lv/wp-content/uploads/2016/10/XIII-BNC.pdf>

Estimation of the influence of K/V on post dialysis urea rebound in HD and HDF with traditional and optical methods

Tomson, Ruth; Fridolin, Ivo; Luman, Merike XIII Baltic Nephrology Conference : October 13-15, 2016, Jūrmala, Latvia : final programme 2016 / p. 2 <http://nefrologs.lv/wp-content/uploads/2016/10/XIII-BNC.pdf>

Free fraction of protein-bound uremic solutes in case of different dialysis modalities

Arund, Jürgen; Tanner, Risto; Luman, Merike; Fridolin, Ivo XIII Baltic Nephrology Conference : October 13-15, 2016, Jūrmala, Latvia : final programme 2016 / p. 17 <http://nefrologs.lv/wp-content/uploads/2016/10/XIII-BNC.pdf>

Total removed beta 2-microglobulin and urea during different dialysis treatment modalities

Holmar, Jana; Luman, Merike; Lauri, Kai; Fridolin, Ivo XIII Baltic Nephrology Conference : October 13-15, 2016, Jūrmala, Latvia : final programme 2016 / p. 21-22 : ill <http://nefrologs.lv/wp-content/uploads/2016/10/XIII-BNC.pdf>