

Metal-binding mechanism of Cox17, a copper chaperone for cytochrome c oxidase

Palumaa, Peep; Kangur, Liina; Voronova, Anastassia; Sillard, Rannar Biochemical journal 2004 / 1, p. 307-314

<https://pubmed.ncbi.nlm.nih.gov/15142040/>

Oxidative switches in functioning of mammalian copper chaperone Cox17

Voronova, Anastassia; Meyer-Klaucke, Wolfram; Meyer, Thomas; Rompel, Anette; Krebs, Bernt; Kazantseva, Jekaterina; Sillard,

Rannar; Palumaa, Peep Biochemical journal 2007 / p. 139-148 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2049083/>

Zn(II) ions co-secreted with insulin suppress inherent amyloidogenic properties of monomeric insulin

Noormägi, Andra; Gavrilova, Julia; Smirnova, Julia; Tõugu, Vello; Palumaa, Peep Biochemical journal 2010 / p. 511-518

<https://pubmed.ncbi.nlm.nih.gov/20632994/>

The first cysteine-rich domain of the receptor GFR[alpha]1 stabilizes the binding of GDNF

Virtanen, Heidi; Yang, Jianmin; Bespalov, Maxim M.; Hiltunen, J.O.; Leppänen, V.M.; Kalkkinen, Nisse; Goldman, A.; Saarma, Mart;

Runeberg-Roos, Pia Biochemical journal 2005 / 3, p. 817-824 <https://pubmed.ncbi.nlm.nih.gov/15610063/>