

Arabidopsis thaliana myosin XIK is involved in root hair as well as trichome morphogenesis on stems and leaves

Ojangu, Eve-Ly; Järve, Kristel; Paves, Heiti; Truve, Erkki Protoplasma 2007 / 3/4, p. 193-202

<https://link.springer.com/article/10.1007/s00709-006-0233-8>

Myosins XI-K, XI-1, and XI-2 are required for development of pavement cells, trichomes, and stigmatic papillae in Arabidopsis

Ojangu, Eve-Ly; Tanner, Krista; Pata, Pille; Järve, Kristel; Holweg, Carola; Truve, Erkki; Paves, Heiti BMC plant biology 2012

<https://pubmed.ncbi.nlm.nih.gov/22672737/>

Myosins XI-K, XI-1, and XI-2 are required for development of pavement cells, trichomes, and stigmatic papillae in Arabidopsis [Online resource]

Ojangu, Eve-Ly; Tanner, Krista; Pata, Pille; Järve, Kristel; Talts, Kristiina; Truve, Erkki; Paves, Heiti Environmental adaptation : from molecules to the planet : the Estonian Centre of Excellence in Environmental Adaptation ENVIRON. Final conference : October 1-3, 2015, Dorpat Conference Centre, Tartu, Estonia : abstract book 2015 / p. 49-50

http://environ.emu.ee/userfiles/environ/Abstract_Book_ENVIRON%20final%20conference%202015.pdf

Redundant functions of class XI myosins of Arabidopsis thaliana

Ojangu, Eve-Ly; Järve, Kristel; Talts, Kristiina; Paves, Heiti; Truve, Erkki 5th EPSO Conference "Plants for Life" : Olos, 29.08.-2.09 2010