

Cu(II) partially protects three histidine residues and the N-terminus of amyloid- β peptide from diethyl pyrocarbonate (DEPC) modification

Friedemann, Merlin; Tõugu, Vello; Palumaa, Peep FEBS Open Bio 2020 / p. 1072-1081 <https://doi.org/10.1002/2211-5463.12857>
[Journal metrics at Scopus](#) [Article at Scopus](#) [Journal metrics at WOS](#) [Article at WOS](#)

Effect of agitation on the peptide fibrillization: Alzheimer's amyloid- b peptide 1-42 but not amylin and insulin fibrils can grow under quiescent conditions

Tiiman, Ann; Noormägi, Andra; Friedemann, Merlin; Krištal, Jekaterina; Palumaa, Peep; Tõugu, Vello Journal of peptide science 2013 / p. 386-391 : ill <https://doi.org/10.1002/psc.2513> [Journal metrics at Scopus](#) [Article at Scopus](#) [Journal metrics at WOS](#) [Article at WOS](#)

Insulin fibrillization at acidic and physiological pH values is controlled by different molecular mechanisms

Noormägi, Andra; Valmsen, Karin; Tõugu, Vello; Palumaa, Peep The protein journal 2015 / p. 398-403 : ill
<https://doi.org/10.1007/s10930-015-9634-x> [Journal metrics at Scopus](#) [Article at Scopus](#) [Journal metrics at WOS](#) [Article at WOS](#)

Interference of low-molecular substances with the thioflavin-T fluorescence assay of amyloid fibrils

Noormägi, Andra; Primar, Kateryna; Tõugu, Vello; Palumaa, Peep Journal of peptide science 2012 / p. 59-64 : ill
<https://pubmed.ncbi.nlm.nih.gov/22083646/>